STANDARD F HDA1C SD BIOSENSOR

STANDARD F HbA1c is a test for quantitaive measurement for glycated hemoglobin (HbA1c) in human capillary venous whole blood. The test is to monitor glycemic control in people with diabetes.

- Optimized performance
- No interference with Hb S, Hb F, Carbamylated Hemoglobin
- Reagents can be stored at room temperature: 2-30°C / 36-86°F
- IFCC / NGSP certified




CE

Specification		
Specimen Type	Whole Blood	
Specimen Volume	5 μl	
Measurement Range	4 - 15 % [NGSP], 20 - 140 mm/mol [IFCC]	
Reference Range	≤ 5.6% (Normal) 5.7 - 6.4% (Prediabetes) ≥ 6.5% (Diabetes) 7% (ADA target for diabetes patients)	
Testing Time	3 mins.	
No interference	Common Hb variants (HbS, HbE, HbC, HbD, HbF)	
Storage Condition	2 - 30°C / 36 - 86°F	

METHODS OF SAMPLE COLLECTION



TEST PROCEDURE





Mix the sample with the extraction buffer and collect all the mixture with the SpoitTM (Blue).



Apply the sample mixture into the sample well of the test device.

METHOD COMPARISON

Reference Method vs STANDARD F HbA1c		
Correlation with HPLC	y=0.9765x + 0.1254, R ² =0.9973, n=80	
CV%	<3%	
Differ (%)	Within 6% (NGSP Criteria)	

ORDERING INFORMATION

Category	Product	Pack Size
Chronic Disease	STANDARD F HbA1c	20 Test
	SDB HbA1c Control	Lv1 x 10 / Lv2 x 10

